

# Effectiveness of statins for secondary prevention in elderly patients after acute myocardial infarction: an evaluation of class effect

**Zheng Zhou, Elham Rahme, Michal Abrahamowicz, Jack V. Tu, Mark J. Eisenberg, Karin Humphries, Peter C. Austin and Louise Pilote**

From the Department of Epidemiology and Biostatistics, McGill University, and the Division of Clinical Epidemiology, Montreal General Hospital, Montréal, Que. (Zhou, Rahme, Abrahamowicz, Pilote); the Institute for Clinical Evaluative Science, Toronto, Ont. (Tu, Austin); the Division of Cardiology, University of British Columbia, and the Centre for Health Evaluation and Outcome Sciences, Vancouver, BC (Humphries); and the Sir Mortimer B. Davis Jewish General Hospital, Montréal, Que. (Eisenberg).

**Correspondence to:** Dr. Louise Pilote, Division of Clinical Epidemiology, Rm. L10-421, Montreal General Hospital, 1650 Cedar Ave., Montréal QC H3G 1A4; fax 514 934-8293; [louise.pilote@mcgill.ca](mailto:louise.pilote@mcgill.ca)

**Background:** Clinical trials have shown the benefits of statins after acute myocardial infarction (AMI). However, it is unclear whether different statins exert a similar effect in reducing the incidence of recurrent AMI and death when used in clinical practice.

**Methods:** We conducted a retrospective cohort study (1997–2002) to compare 5 statins using data from medical administrative databases in 3 provinces (Quebec, Ontario and British Columbia). We included patients aged 65 years and over who were discharged alive after their first AMI-related hospital stay and who began statin treatment within 90 days after discharge. The primary end point was the combined outcome of recurrent AMI or death from any cause. The secondary end point was death from any cause. Adjusted hazard ratios (HRs) for each statin compared with atorvastatin as the reference drug were estimated using Cox proportional hazards regression analysis.

**Results:** A total of 18 637 patients were prescribed atorvastatin ( $n = 6420$ ), pravastatin ( $n = 4480$ ), simvastatin ( $n = 5518$ ), lovastatin ( $n = 1736$ ) or fluvastatin ( $n = 483$ ). Users of different statins showed similar baseline characteristics and patterns of statin use. The adjusted HRs (and 95% confidence intervals) for the combined outcome of AMI or death showed that each statin had similar effects when compared with atorvastatin: pravastatin 1.00 (0.90–1.11), simvastatin 1.01 (0.91–1.12), lovastatin 1.09 (0.95–1.24) and fluvastatin 1.01 (0.80–1.27). The results did not change when death alone was the end point, nor did they change after adjustment for initial daily dose or after censoring of patients who switched or stopped the initial statin treatment.

**Interpretation:** Our results suggest that, under current usage, statins are equally effective for secondary prevention in elderly patients after AMI.